

# Individual heterogeneity and identifiability in capture–recapture models

W. A. Link

Link, W. A., 2004. Individual heterogeneity and identifiability in capture–recapture models. *Animal Biodiversity and Conservation*, 27.1: 87–91.

## Abstract

*Individual heterogeneity and identifiability in capture–recapture models.*— Individual heterogeneity in detection probabilities is a far more serious problem for capture–recapture modeling than has previously been recognized. In this note, I illustrate that population size is not an identifiable parameter under the general closed population mark–recapture model  $M_h$ . The problem of identifiability is obvious if the population includes individuals with  $p_i = 0$ , but persists even when it is assumed that individual detection probabilities are bounded away from zero. Identifiability may be attained within parametric families of distributions for  $p_i$ , but not among parametric families of distributions. Consequently, in the presence of individual heterogeneity in detection probability, capture–recapture analysis is strongly model dependent.

Key words: Capture–recapture, Detection probability, Heterogeneity, Identifiability, Population estimation.

## Resumen

*Heterogeneidad individual e identificabilidad en modelos de captura–recaptura.*— La heterogeneidad individual en las probabilidades de detección representa un problema para la modelación del procedimiento de captura–recaptura mucho más serio de lo que previamente se había reconocido. En este artículo se demuestra que el tamaño de la población no constituye un parámetro identificable en el modelo general  $M_h$  que emplea técnicas de marcaje–recaptura de poblaciones cerradas. El problema de la identificabilidad resulta evidente si la población incluye individuos con  $p_i = 0$ , pero sigue persistiendo aun cuando se presuponga que las probabilidades de detección individual se han alejado de cero. La identificabilidad puede conseguirse en familias paramétricas de distribuciones para  $p_i$ , pero no entre familias paramétricas de distribuciones. Por consiguiente, si se da una heterogeneidad individual en la probabilidad de detección, el análisis de captura–recaptura depende considerablemente del modelo considerado.

Palabras clave: Captura–recaptura, Probabilidades de detección, Heterogeneidad, Identificabilidad, Estimación de la población.

William A. Link, USGS Patuxent Wildlife Research Center, 11510 American Holly Drive, Laurel, Maryland 20708 U.S.A.

## Introduction

Let  $X_i$ ,  $i = 1, 2, \dots, N$  be independent binomial random variables, with common index  $T$  and success rates  $p_i$ , sampled independently from distribution  $g(p)$ . Further, let

$$f_j = \sum_{i=1}^N 1(X_i = j) \quad j = 0, 1, \dots, T$$

where  $1(\cdot)$  is the indicator function. Having observed  $f^c = (f_1, f_2, \dots, f_T)$ , the problem is to estimate  $N$ , or equivalently, to predict  $f_0$ . This is the closed population capture–recapture model  $M_h$ :  $N$  is the unknown population size,  $X_i$  is the number of times animal  $i$  is captured in  $T$  sampling occasions,  $f_j$  is the number of animals captured exactly  $j$  times.

Numerous methods for estimating  $N$  exist, ranging from the jackknife method of Burnham & Overton (1978), to finite mixture models (Norris & Pollock, 1996; Pledger, 2000), and including parametric models such as the logit normal (Coull & Agresti, 1999) and beta models (Dorazio & Royle, 2003). Given the restrictions on  $g(p)$  implicit to these methods, estimation of  $N$  is usually successful. However, as will be demonstrated here and elsewhere (Link, 2003),  $N$  is not identifiable without untestable model assumptions restricting the set of distributions  $g(p)$ .

One example of this difficulty is well known. If the population consists of  $N_1$  individuals with  $p_i = 0$ , and  $N_2$  individuals with  $p_i > 0$ , an analyst of model  $M_h$  can at best estimate  $N_2$ , rather than  $N = N_1 + N_2$ . This circumstance is generally dismissed with the assertion that "we're only estimating the observable portion of the population."

But what of animals with low but nonzero detection probabilities? These are clearly the ones which present the challenge to capture–recapture analysis. Huggins (2001), seeking to identify restrictions on the collection of distributions that would ensure identifiability, focused his attention on removing difficulties associated with low detection probabilities. The condition he considered was that  $g(p)$  places no mass on values of  $p < 1 - (1 - \gamma)^{1/T}$  for a fixed value  $\gamma \in (0, 1)$ . This means that every individual has probability of at least  $\gamma > 0$  of being captured on one of the  $T$  sampling occasions. Huggins concluded that if the converse of his Theorem 3 were true (he describes this as "difficult to establish" and "an open question") then the condition would be sufficient to ensure identifiability.

The restriction is not sufficient, as is demonstrated by example 1, below. It is possible to construct 2 distinct distributions,  $g_1(p) \neq g_2(p)$ , each with support bounded away from zero, the two distributions producing identical sampling distributions for the observed data  $f^c$ , but leading to contradictory inference about  $f_0$ .

Stronger restrictions, or at least *different* restrictions are required, to ensure identifiability of  $N$ . For example, Burnham's (1972) thesis includes a demonstration that restricting attention to beta distrib-

uted heterogeneity leads to identifiability of  $N$ . Thus we can feel confident dealing with model  $M_h$  if we are confident that  $g(p)$  is a beta distribution. But what if, unbeknownst to us,  $g(p)$  is a logit normal distribution? It can be demonstrated by example that the sampling distribution of  $f^c$  induced by a beta distribution can be very closely approximated by the sampling distribution of  $f^c$  induced by a logit normal distribution, but with substantially different inferences about  $N$ . (See example 2, below.) The inferences are distinct, but there is no way, on the basis of data  $f^c$  to decide which is correct (except with vast sample sizes). Since it is unlikely that one will have epistemological grounds for assuming the beta distribution over the logit normal (or other distributions, such as the log gamma; see Link, 2003), it seems faint comfort to learn that  $N$  is identifiable within any one of these classes.

My third example, below, shows that if nature is perverse in its selection of  $g(p)$ , the sampling distribution of  $f^c$  can be strongly and misleadingly suggestive of a particular form for  $g(p)$ , even for a variety of values for  $T$ .

## Additional notation

Let  $n$  denote the number of distinct animals ever sighted, i.e.,  $n = f_1 + f_2 + \dots + f_T$ . I refer to the data  $f^c$  as the observed frequency distribution, and to  $f = (f_0, f_1, \dots, f_T)$  as the complete frequency distribution. The vectors  $f$  and  $f^c$  are multinomial random variables with indices  $N$  and  $n$ , respectively, and  $j$ th cell probabilities designated by  $\pi(j)$  and  $\pi^c(j)$ , respectively. These are related by  $\pi^c(j) = \pi(j)/1 - \pi(0)$ . Under model  $M_h$ , we have

$$\pi(j) = \int_0^1 \binom{T}{j} p^j (1-p)^{T-j} g(p) dp \quad (1)$$

Substituting an estimate  $\hat{g}(p)$  for  $g(p)$  in (1), one obtains estimates  $\hat{\pi}(j)$ ,  $j = 0, 1, 2, \dots, n$ .

It is easily verified that

$$E(f_0) = \frac{\pi(0)}{1 - \pi(0)} E(n) \quad (2)$$

Thus, it is natural to predict the number of individuals not seen by

$$\hat{f}_0 = \frac{\hat{\pi}(0)}{1 - \hat{\pi}(0)} n$$

and to predict the unknown population size by  $\hat{N} = n + \hat{f}_0$

## Example 1

Suppose that  $T = 6$ , that  $g_1(p)$  corresponds to a uniform distribution on (0.008512, 0.76), and that  $g_2(p)$  corresponds to a 3–point mixture placing masses {0.350739, 0.414090, 0.235172} on values {0.161937, 0.449089, 0.692734}. The minimum value of  $p$  attainable under either model is 0.008512 =  $1 - (1 - \gamma)^{1/T}$ , for  $\gamma = 0.050$ ; every animal has at

Table 1. Cell probabilities for  $\pi(x)$  and  $\pi^c(x)$  for complete and observed frequency distributions of Example 1.

Tabla 1. Probabilidades de cada celda para  $\pi(x)$  y  $\pi^c(x)$  de las frecuencias de distribución completas y observadas del Ejemplo 1.

		0	1	2	3	4	5	6
Unif (a,b)	$\pi(x)$	.179048	.189616	.188057	.178371	.147685	.089382	.027840
	$\pi^c(x)$	–	.230971	.229072	.217273	.179895	.108876	.033912
3 pt. mixture	$\pi(x)$	.133294	.200184	.198538	.188312	.155916	.094364	.029392
	$\pi^c(x)$	–	.230971	.229072	.217273	.179895	.108876	.033912

least a 5% chance of being caught on one or more sampling occasions.

The cell probabilities for  $f$  and  $f^c$  are given in table 1. Note that the sampling distribution of the data  $f^c$  is identical for the two distributions  $g(p)$ , but that the predicted value of  $f_0$  is nearly half again as large under the uniform distribution as under the two-point mixture: with  $n = 100$ , the prediction of  $f_0$  under the uniform specification is  $100(0.179)/(1 - 0.179) \approx 22$ , while the prediction of  $f_0$  under the 3-point specification is  $100(0.133)/(1 - 0.133) \approx 15$ .

I describe the method used for constructing Example 1 in presenting Example 3, below. Example 1 may be of special interest to analysts, since the two distributions correspond to models that could be fit based on observations from  $T = 6$  sampling periods. It is worth mentioning that the problem of identifiability does not depend on both models being fittable, a point to which I return in presenting Example 3.

### Example 2

Let  $T = 5$ , and  $g_1(p)$  represent the distribution resulting from the assumption that  $\text{logit}(p)$  has a normal distribution with mean of  $-1.75$  and standard deviation of  $2.00$ . I calculated the sampling distribution for  $f$  and  $f^c$  by numeric integration over a grid of  $100,000$  points.

Next, I minimized the Kullback–Leibler (KL) distance from the distribution of  $f^c$  induced by  $g_1(p)$  to the distribution of  $f^c$  induced by  $g(p)$  in the beta family of distributions (for details on the KL–distance; see Agresti, 1990: p. 241). The resulting beta distribution has parameters  $a = 0.2512$  and  $b = 1.1300$ . This beta distribution and the logit normal distribution described in the previous paragraph are plotted in figure 1.

The observed and complete frequency distributions are given in table 2. Note that while the observed frequency distributions are not identical, they are close enough to be virtually indistinguish-

able except with extremely large samples. The discrepancy in predictions of  $f_0$  is substantial: based on  $n = 100$ , the predictions are 83 (for the logit-normal model) and 156 (for the beta model).

### Example 3

Expanding the term  $(1 - p)^{T-j}$  in equation (1) by means of the binomial theorem, it is seen that the values  $\pi(x)$  are linear combinations of the first  $T$  moments of distribution  $g(p)$ , for  $x = 1, 2, \dots, T$ . The same is true for

$$1 - \pi(0) = \sum_{x=1}^T \pi(x)$$

Let  $m_g(j)$  denote the  $j$ th moment of  $g(p)$ . If we could construct a distribution  $h(p)$  with moments  $m_h(j) = cm_g(j)$ , for some  $c \neq 1$ , and for  $j = 1, 2, \dots, T$ , the observations in the previous paragraph, and the relation  $\pi^c(j) = \pi(j)/(1 - \pi(0))$  lead to the conclusion that distributions  $g(p)$  and  $h(p)$  will induce the same values  $\pi^c(j)$ , but different values for  $\pi(0)$ . Using subscripts  $g$  and  $h$  to distinguish the values of  $\pi(0)$ , we obtain the relation

$$\pi_h(0) = (1 - c) + c\pi_g(0) \quad (3)$$

Thus these distinct distributions of heterogeneity will lead to the same sampling distributions for observed data, but different predictions for  $f_0$ , and consequently, for  $N$ .

If  $g(p)$  is the uniform distribution,  $m_g(j) = 1/(j + 1)$ . Consider the distribution function  $h(p) = 1/15 (114 - 4950p + 79200p^2 - 600600p^3 + 2522520p^4 - 6306300p^5 + 9609600p^6 - 8751600p^7 + 4375800p^8 - 923780p^9)$ .

This distribution is plotted along with the uniform distribution in figure 2. Straightforward calculation shows that the moments of  $h(p)$  are  $m_h(j) = c/(j + 1)$ , with  $c = 14/15$ , for  $j = 1, 2, \dots, 9$ .

Thus for a study involving any number of sampling occasions up to  $T = 9$ , the data produced with heterogeneity distribution  $h(p)$  will be




Fig. 1. Density functions used in Example 2. Dashed line represents beta distribution with parameters  $a = 0.2512$  and  $b = 1.1300$ ; solid line is distribution of  $p$  corresponding to  $\text{logit}(p)$  having a normal distribution with mean of  $-1.75$  and standard deviation of  $2.00$ . Note that  $x$ -axis has been distorted to accentuate the differences between the two densities.

Fig. 1. Funciones de densidad utilizadas en el Ejemplo 2. La línea discontinua representa la distribución beta con los parámetros  $a = 0,2512$  y  $b = 1,300$ ; la línea continua es la distribución de  $p$  correspondiente al  $\text{logit}(p)$  que presenta una distribución normal con una media de  $-1,75$  y una desviación estándar de  $2,00$ . Nótese que el eje  $x$  se ha distorsionado para acentuar las diferencias entre las dos densidades.

Table 2. Cell probabilities for  $\pi(x)$  and  $\pi^C(x)$  for complete and observed frequency distributions of Example 2.

Tabla 2. Probabilidades de cada celda para  $\pi(x)$  y  $\pi^C(x)$  de las frecuencias de distribución completas y observadas del Ejemplo 2.

		0	1	2	3	4	5
Logit-normal	$\pi(x)$	0.454	0.208	0.126	0.090	0.070	0.052
	$\pi^C(x)$	–	0.381	0.231	0.165	0.128	0.095
Beta	$\pi(x)$	0.609	0.149	0.090	0.065	0.050	0.037
	$\pi^C(x)$	–	0.381	0.231	0.166	0.127	0.095

indistinguishable from data produced under a uniform distribution of heterogeneity. However, the predictions of  $f_0$  will differ substantially. Since under the uniform distribution  $\pi_g(0) = 1/(T + 1)$ , we find from (3) and (2) that the prediction of  $f_0$  based on the uniform distribution will be smaller than the prediction based on  $h(p)$  by a factor of  $14 / (T + 15)$ .

Some might dismiss this example on the grounds that they would never even consider fitting a distribution that looks like  $h(p)$ ; to this, I reply "That's exactly my point!" If nature perversely selects  $h(p)$  as the distribution of heterogeneity, and we are (excusably) misled into assuming a uniform distri-

bution, our predictions of  $f_0$  will be too small by a factor of  $14 / (T + 15)$ , and we'll never know the difference.

### Conclusions and discussion

Population size  $N$  is not an identifiable parameter under model  $M_h$ , except under the imposition of untestable model assumptions. Thus estimation of population size, in the presence of individual heterogeneity in detection is inevitably model based, much the same as the analysis of oft-reviled count survey data.




Fig. 2. Density functions used in Example 3. The first nine moments of the nonuniform density are precisely 14/15th's the size of the corresponding moments of the uniform density.

Fig. 2. Funciones de densidad utilizadas en el Ejemplo 3. Los primeros nueve momentos de la densidad no uniforme equivalen precisamente a 14/15 del tamaño de los momentos homólogos de la densidad uniforme.

It is worth considering what the further implications of this finding are, particularly for open population models. Some early indications (Link, 2003) are that while estimates of population sizes may be biased in a manner similar to that described here for closed population estimation, survival estimates may be less sensitive to heterogeneity in detection rates. On the other hand, since capture–mark–recapture experiments essentially create "populations" of marked animals that are closed except to mortality, it is possible that time variation in detection rates might induce bias in survival estimates.

The problems presented here should come as no surprise. Indeed, without specific parametric models for the heterogeneity in  $p$ , we find ourselves in the unpleasant circumstances described in the classic paper of Kiefer & Wolfowitz (1956) which demonstrated, among other things, that maximum likelihood estimates of parameters of interest may be asymptotically biased, and badly so, if the number of nuisance parameters is allowed to increase without bound. This is precisely the situation under model  $M_h$ , with individual detection probabilities in the role of nuisance parameters.

#### References

- Agresti, A., 1990. *Categorical Data Analysis*. New York, Wiley.
- Burnham, K. P., 1972. Estimation of population size in multiple capture–recapture studies when capture probabilities vary among animals. Ph. D. Thesis, Oregon State Univ., Corvallis.
- Burnham, K. P. & Overton, W. S., 1978. Estimation of the size of a closed population when capture probabilities vary among animals. *Biometrika*, 65: 625–633.
- Coull, B. A. & Agresti, A., 1999. The use of mixed logit models to reflect heterogeneity in capture–recapture studies. *Biometrics*, 55: 294–301.
- Dorazio, R. M. & Royle, J. A., 2003. Mixture models for estimating the size of a closed population when capture rates vary among individuals. *Biometrics*, 59: 351–364.
- Huggins, R., 2001. A note on the difficulties associated with the analysis of capture–recapture experiments with heterogeneous capture probabilities. *Statistics and Probability Letters*, 54: 147–152.
- Kiefer, J. & Wolfowitz, J., 1956. Consistency of the maximum likelihood estimator in the presence of infinitely many nuisance parameters. *Annals of Mathematical Statistics*, 27: 887–906.
- Link, W. A., 2003. Nonidentifiability of population size from capture–recapture data with heterogeneous detection probabilities. *Biometrics*, 59: 1125–1132.
- Norris, J. L. & Pollock, K. H., 1996. Nonparametric MLE under two closed–capture models with heterogeneity. *Biometrics*, 52: 639–649.
- Pledger, S., 2000. Unified maximum likelihood estimates for closed capture–recapture models using mixtures. *Biometrics*, 56: 434–442.